

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: GUPDOCADT

(Page 1 of 1)

| DOCTOR'S ORDERS | Ht | cm | Wt | kg | BSA | m² |
|--|-----------------|-----------|----------|---------|---------------------------------------|----|
| REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form | | | | | | |
| DATE: To be | e given: | | | Cycle # | of 6 | ; |
| Date of Previous Cycle: | | | | | | |
| Delay treatment week(s) | | | | | | |
| CBC & Diff, platelets day of treatment | | | | | | |
| May proceed with doses as written if within 96 hours ANC greater than or equal to 1.5 x 10 ⁹ /L, platelets greater than | | | | | | |
| 90 x 10 ⁹ /L (for cycles 1 and 4, also if within 96 hours total bilirubin less than or equal to ULN, alkaline phosphatase less than 2.5 x ULN (unless bone metastases), and AST and/or ALT less than or equal to 1.5 x ULN) | | | | | | |
| Dose modification for: Hematology Other Toxicity: | | | | | | |
| Proceed with treatment based on blood wor | | | | | · · · · · · · · · · · · · · · · · · · | |
| | | | | | | |
| PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm | | | | | | |
| dexamethasone 8 mg PO bid for 3 days, starting one day prior to treatment; patient must receive a minimum of 3 doses pretreatment | | | | | | |
| Optional: Frozen gloves starting 15 minutes before DOCEtaxel infusion until 15 minutes after end of DOCEtaxel infusion; gloves should be changed after 45 minutes of wearing. | | | | | | |
| ☐Other: | | | | | | |
| **Have Hypersensitivity Reaction Tray and Protocol Available** | | | | | | |
| TREATMENT: | | | | | | |
| DOCEtaxel 75 mg/m ² x BSA = mg | | | | | | |
| Dose Modification:% = mg/m ² x BSA = mg | | | | | | |
| IV in 250 to 500 mL (non-DEHP bag) NS over one hour (use non-DEHP tubing) | | | | | | |
| | | | | | | |
| Remember to commence standard androgen deprivation therapy (i.e. LHRH agonist, LHRH antagonist, anti-androgen) | | | | | | |
| | | | | | | |
| RETURN APPOINTMENT ORDERS | | | | | | |
| Return in <u>three</u> weeks for Doctor and Cycle | e | | | | | |
| Last Cycle. Return in week(s). | | | | | | |
| CBC & Diff, platelets and PSA prior to each of | cycle | | | | | |
| Prior to Cycle 4 and as clinically indicated: A bilirubin, LDH | ALT, alkaline p | hosphatas | e, total | | | |
| ☐Other tests: | | | | | | |
| Consults: | | | | | | |
| | | | | | | |
| See general orders sheet for additional re | equests. | | | | | |
| DOCTOR'S SIGNATURE: | | | | SIG | BNATURE | • |
| | | | | UC | | |