

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: GOOVCAG

Page 1 of 1

DOCTOR'S ORDERS Htcm Wtkg Bit	SAm²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form	
DATE: To be given: Cycle #	<i>‡</i> :
Date of Previous Cycle:	
 Delay treatment week(s) CBC & Diff, Platelets day of treatment May proceed with doses as written if within 24 hours ANC greater than or equal to 1.0 x 10⁹/L, platelets greater than or equal to 100 x 10⁹/L Dose modification for: Hematology Other Toxicity Proceed with treatment based on blood work from 	
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm	
dexamethasone 8 mg PO 30 to 60 minutes prior to CARBOplatin	
AND select I ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin	
ONE of the aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and	
following: Ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin	
netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to CARBOplatin	
If additional antiemetic required:	
□ OLANZapine □ 2.5 mg or □ 5 mg or □ 10 mg (select one) PO 30 to 60 minutes prior to CARBOplatin	
Other:	
Have Hypersensitivity Reaction Tray and Protocol Available CHEMOTHERAPY:	
DAY 1 gemcitabine \square 800 mg/m ² OR \square mg/m ² (select one) x BSA =mg \square Dose Modification:% =mg/m ² x BSA =mg IV in 250 mL NS over 30 minutes. (Maximum Dose = 2000 mg) CARBOplatin AUC \square 5 OR \square 6 OR \square 4 (select one) X (GFR+25) =mg \square Dose Modification:% =mg IV in 100 to 250 mL NS over 30 minutes, after gemcitabine.	
DAY 8 gemcitabine	
RETURN APPOINTMENT ORDERS	
 Return in <u>three</u> weeks for Doctor and Cycle # Book chemo Day 1 and 8 Last Cycle. Return in week(s). 	
 CBC & Diff, platelets, creatinine, prior to Day 1 each cycle CBC & Diff, platelets, prior to Day 8 each cycle Other tests: Nuclear renogram for GFR If Clinically Indicated: CA -125 CA 15-3 CA 19-9 prior to each cycle Consults: See general orders sheet for additional requests. 	
DOCTOR'S SIGNATURE:	SIGNATURE:
	UC: