PROTOCOL CODE: LUAJNIVPP

DOCTOR'S ORDERS

Ht _____**cm** **Wt** _____**kg** **BSA** _____**m²**

REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form

DATE: **To be given:** **Cycle #:**

Date of Previous Cycle:

☐ Delay treatment _____ week(s)
☐ **CBC & Diff, platelets** day of treatment

May proceed with pemetrexed, CISplatin, CARBOplatin as written if within 96 hours **ANC greater than or equal to 1.5 x 10^9/L, platelets greater than or equal to 100 x 10^9/L, and creatinine clearance greater than or equal to 45 mL/minute (**for pemetrexed and CARBOplatin**),** or **greater than or equal to 60 mL/minute** (for CISplatin)

May proceed with nivolumab if within 96 hours **creatinine less than or equal to 1.5 times the upper limit of normal** *and* **less than or equal to 1.5 times the baseline, ALT less than or equal to 3 times the upper limit of normal, total bilirubin less than or equal to 1.5 times the upper limit of normal**

Dose modification for: ☐ **Hematology** ☐ **Other Toxicity:** _____
Proceed with treatment based on blood work from _____

PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm _____.

dexamethasone ☐ **8 mg** or ☐ **12 mg** (select one) PO 30 to 60 minutes prior to treatment

AND select **ONE** of the following:	☐	**aprepitant 125 mg** PO 30 to 60 minutes prior to treatment, and **ondansetron 8 mg** PO 30 to 60 minutes prior to treatment
	☐	**netupitant-palonosetron 300 mg-0.5 mg** PO 30 to 60 minutes prior to treatment
	☐	**ondansetron 8 mg** PO 30 to 60 minutes prior to treatment

If additional antiemetic required:
☐ **OLANZapine** ☐ **2.5 mg** or ☐ **5 mg** or ☐ **10 mg** (select one) PO 30 to 60 minutes prior to treatment

Ensure patient is taking **folic acid** and has had **vitamin B12** injection starting at least 7 days prior to first cycle, and to continue while on treatment, until 21 days after last pemetrexed dose.

For prior infusion reaction:
☐ **diphenhydrAMINE 50 mg** PO 30 minutes prior to treatment
☐ **acetaminophen 325 to 975 mg** PO 30 minutes prior to treatment
☐ **hydrocortisone 25 mg** IV 30 minutes prior to treatment
☐ **Other:**

****Have Hypersensitivity Reaction Tray & Protocol Available****

PREHYDRATION:

1000 mL NS over 1 hour prior to CISplatin

Continued on page 2

DOCTOR'S SIGNATURE: **SIGNATURE:**

UC:

PROTOCOL CODE: LUAJNIVPP

DATE:

****Have Hypersensitivity Reaction Tray & Protocol Available****

CHEMOTHERAPY:

nivolumab 4.5 mg/kg x _____kg = _____ mg **(max. 360 mg)**
IV in 50 to 100 mL NS over 30 minutes using a 0.2 micron in-line filter (may be given during prehydration)

pemetrexed 500 mg/m^2 x BSA = _____ mg
☐ Dose Modification: _____% = _____ mg/m^2 x BSA = _____ mg
IV in 100 mL NS over 10 minutes (may be given during prehydration)

Select one:
☐ **CISplatin 75 mg/m^2** x BSA = _____ mg
☐ Dose Modification: _____% = _____ mg/m^2 x BSA = _____ mg
IV in 500 mL NS, with potassium chloride 20 mEq, magnesium sulphate 1 g and mannitol 30 g over 1 hour
OR
☐ **CARBOplatin AUC 5 x (GFR + 25)** = _____ mg IV in 100 to 250 mL NS over 30 minutes

RETURN APPOINTMENT ORDERS

☐ Return in **three** weeks for Doctor and Cycle _____.
☐ Last Cycle. Return in _____ week(s).

CBC & Diff, platelets, creatinine, alkaline phosphatase, ALT, total bilirubin, LDH, sodium, potassium, TSH, creatine kinase, random glucose prior to each treatment

CBC & Diff, platelets weekly during Cycles 1 and 2

Vitamin B12 injection required every 9 weeks. Patient to obtain supply.
☐ This patient to receive injection in clinic. Next injection due by _____.

If clinically indicated: ☐ **ECG** ☐ **Chest X-ray**

☐ **serum HCG** or ☐ **urine HCG** (select one) – required for woman of child bearing potential

☐ **Free T3 and free T4** ☐ **lipase** ☐ **morning serum cortisol**

☐ **serum ACTH levels** ☐ **testosterone** ☐ **estradiol** ☐ **FSH** ☐ **LH**

☐ **Weekly nursing assessment**

☐ **Other consults**

☐ **See general orders sheet for additional requests.**

DOCTOR'S SIGNATURE:

SIGNATURE:

UC: