

**PROTOCOL CODE: SMAVIPNI
(Induction)**

Page 1 of 1

DOCTOR'S ORDERS		Wt _____ kg
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form		
DATE:	To be given:	Cycle #:
Date of Previous Cycle: _____		
<input type="checkbox"/> Delay treatment _____ week(s) <input type="checkbox"/> Delay for toxicity Type of toxicity _____		
May proceed with doses as written if within 96 hours ALT less than or equal to 3 times the upper limit of normal, bilirubin less than or equal to 1.5 times the upper limit of normal, creatinine less than or equal to 1.5 times the upper limit of normal and less than or equal to 1.5 X baseline. Proceed with treatment based on blood work from _____		
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm _____. For prior infusion reaction: <ul style="list-style-type: none"> <input type="checkbox"/> diphenhydrAMINE 50 mg PO 30 minutes prior to treatment <input type="checkbox"/> acetaminophen 325 to 975 mg PO 30 minutes prior to treatment <input type="checkbox"/> hydrocortisone 25 mg IV 30 minutes prior to treatment 		
IMMUNOTHERAPY: nivolumab 1 mg/kg x _____ kg = _____ mg every 3 weeks IV in 25 to 50 mL NS over 30 minutes using a 0.2 micron in-line filter* ipilimumab 3 mg/kg x _____ kg = _____ mg every 3 weeks IV in 50 to 250 mL NS over 30 minutes using a 0.2 micron in-line filter* * Use separate infusion line and filter for each drug		
RETURN APPOINTMENT ORDERS		
<input type="checkbox"/> Return in three weeks for Doctor and Cycle _____. (Cycle 5 maint. phase nivolumab q2wk) <input type="checkbox"/> Return in six weeks for Doctor and Cycle 5. (maint. phase nivolumab for q4wk dose only)		
CBC and diff, platelets, creatinine, alkaline phosphatase, ALT, total bilirubin, LDH, sodium, potassium, creatine kinase (CK), TSH, glucose prior to each treatment Weekly nursing assessment If clinically indicated: <input type="checkbox"/> ECG <input type="checkbox"/> Chest X-ray <input type="checkbox"/> serum HCG or <input type="checkbox"/> urine HCG – required for woman of child bearing potential <input type="checkbox"/> Free T3 and free T4 <input type="checkbox"/> lipase <input type="checkbox"/> morning serum cortisol <input type="checkbox"/> serum ACTH levels <input type="checkbox"/> testosterone <input type="checkbox"/> estradiol <input type="checkbox"/> FSH <input type="checkbox"/> LH <input type="checkbox"/> Other consults: <input type="checkbox"/> See general orders sheet for additional requests.		
DOCTOR'S SIGNATURE:		SIGNATURE:
		UC: