

Summer 2011 SON Newsletter References

Improving the Hereditary Cancer Program Referral Process (pg. 2)

1. Armel, S. R., McCuaig, J., Finch, A., Demsky, R., Panzarella, T., Murphy, J., et al. The effectiveness of family history questionnaires in cancer genetic counseling. *Journal of Genetic Counseling* 2009; 18 (4), 366-378.
2. O'Neil, S. M., Peters, J. A., Vogel, V. G., Feingold, E., Rubinstein, W. S. Referral to cancer genetic counseling: Are there stages of readiness? *American Journal of Medical Genetics* 2006; 142C, 221-231.
3. Schlich-Bakker, K. J., ten Kroode, H. F. J., Warmlam-Rodenhuis, C. C., van den Bout, J., & Ausems, M. G. E. M. Barriers to participating in genetic counseling and BRCA testing during primary treatment for breast cancer. *Genetics In Medicine* 2007; 9 (11), 766-777.

'Ovarian' Cancer Prevention: Simple Salpingectomy (pg. 3)

1. Hogg R, Friedlander M. Biology of epithelial ovarian cancer: implications for screening women at high genetic risk. *J Clin Oncol* 2004; 22:1315–1327.
2. Auersperg N, Wong AS, Choi KC, et al. Ovarian surface epithelium: biology, endocrinology, and pathology. *Endocr Rev* 2001; 22:255–288.
3. Shaw P, Rouzbahman M, Murphy J, et al. Mucosal epithelial proliferation and p53 over expression are frequent in women with BRCA mutations (Abstract). *Mod Pathol* 2004; 17(Supp 1):214A.
4. Crum CP, Drapkin R, Kindelberger D, et al. Lessons from BRCA: the tubal fimbria emerges as an origin for pelvic serous cancer. *Clin Med Res* 2007; 5:35–44.
5. Lee Y, Miron A, Drapkin R, et al. A candidate precursor to serous carcinoma that originates in the distal fallopian tube. *J Pathol* 2007; 211:26–35.
6. Medeiros F, Muto MG, Lee Y, et al. The tubal fimbria is a preferred site for early adenocarcinoma in women with familial ovarian cancer syndrome. *Am J Surg Pathol* 2006; 30:230–236.
7. Lee Y, Miron A, Drapkin R, Nucci MR, Medeiros F, Saleemuddin A, et al. A candidate precursor to serous carcinoma that originates in the distal fallopian tube. *J Pathol* 2007; 211(1):26-35.
8. Crum CP, Drapkin R, Kindelberger D, Medeiros F, Miron A, Lee Y. Lessons from BRCA: the tubal fimbria emerges as an origin for pelvic serous cancer. *Clin Med Res* 2007; 5(1):35-44.

9. Salvador S, Gilks B, Köbel M , Huntsman D, Rosen B, Miller D. The fallopian tube: primary site of most pelvic high-grade serous carcinomas. *Int J Gynecol Ca* 2009; 19:58-64.
10. Karst AM, Levanon K, Drapkin R . Modeling high-grade serous ovarian carcinogenesis from the fallopian tube. *Proc Natl Acad Sci USA*. 2011 May 3; 108(18):7547-52.
11. CIHI Health Indicators 2010:
http://secure.cihi.ca/cihilweb/products/Healthindicators2010_md_en.pdf
12. Rebbeck TR, Lynch HT, Neuhausen SL, Narod SA, Van't Veer L, Garber JE et al. *N Engl J Med* 2002. 346: 1616–1622.
13. Rutter JL , Wacholder S , Chetrit A , et al . Gynecologic surgeries and risk of ovarian cancer in women with BRCA1 and BRCA2 Ashkenazi founder mutations: an Israeli population-based case-control study . *J Natl Cancer Inst*. 2003; 95 (14): 1072 – 1078 .
14. Domchek SM, Friebel TM, Neuhausen SL, Wagner T, Evans G, Isaacs C et al. (2006). *Lancet Oncol* 2006; 7: 223–229.
15. Finch A et al. Salpingo-oophorectomy and the risk of ovarian, fallopian tube, and peritoneal cancers in women with a BRCA1 or BRCA2 mutation. *JAMA* 2006; 296: 185-192.

Supporting Chinese-speaking cancer patients to cope with a cancer diagnosis and access to support (pgs. 6&7)

1. Bell K, Lee J, Ristovski-Slijepcevic S. Perceptions of food and eating amongst Chinese cancer patients: Findings of an ethnographic study. *Cancer Nursing* 2009; 32(2):1-9.
2. Bell K, Lee J. (2009). Chinese cancer support group study: Key findings. Available at <http://www.bccancer.bc.ca/NR/rdonlyres/E6F649B9-761C-4C51-89E0-C2F0834B8DCC/32818/FINALCCSGREPORT9March.pdf>
3. Barroetavena, MC., Kwong, S., Chiu, L., Doll, R., The, C., Chan, T. (2006). Psychosocial Needs of Chinese Cancer Patients – Are We Providing Cultural Competent Care Services. CAPO Poster.