

For the Patient: **BRAJACT**

Other names: Neoadjuvant or Adjuvant Therapy for Breast Cancer using DOXOrubicin and Cyclophosphamide followed by PACLitaxel

BR = **BR**east

AJ = **AdJ**uvant

A = **DOXO**rubicin (**ADRIAMYCIN**®)

C = **C**yclophosphamide

T = **PAC**Litaxel (**TAXOL**®)

Uses:

- BRAJACT is a drug treatment given after breast cancer **diagnosis** in the hope of destroying breast cancer cells that may have spread to other parts of your body. **It can be given before (called neoadjuvant) or after (called adjuvant) breast cancer surgery.** This treatment may reduce the chance of your breast cancer coming back and may improve your overall survival or chance of a cure.

Treatment Plan:

- Your treatment plan consists of 8 chemotherapy cycles. Total time on drug treatment would be up to about 6 months. A cycle length is 3 weeks. All the drugs are given intravenously at every visit. For each cycle, you will need to have a blood test and see your oncologist before the treatment. The dose and timing of your chemotherapy may be changed based on your blood counts and/or other side effects.
- Doxorubicin and Cyclophosphamide (AC) are given for the first 4 cycles. Each treatment takes about one hour. You will be given a prescription for anti-nausea drugs to take 30 minutes before the treatment and again about 6-8 hours after, and will usually be on anti-nausea drugs the following 2 days also.
- The next 4 cycles with Paclitaxel take about 4 hours in the chemotherapy room. You will be given some intravenous medications before the Paclitaxel to prevent or lessen the chance of allergy.
- If needed, radiation therapy will start after your 8th chemotherapy cycle is finished and hormone treatments may also be started, if your oncologist has recommended these for you.

Instructions:

- Bring your anti-nausea drugs with you to take before each AC treatment. You will also need to take your anti-nausea drugs at home after therapy. It is easier to prevent nausea than treat it once it has occurred, so follow directions closely.
- Drink lots of fluids for the first day or two after chemotherapy, especially while on AC (8 or more cups a day).
- Check with your doctor or pharmacist before you start taking any new drugs. Other drugs such as Phenytoin (DILANTIN®), Warfarin (COUMADIN®), Digoxin (LANOXIN®), and Thiazide diuretics (“water pills”) may interact with BRAJACT.
- You may drink small amounts of alcohol, as it will not affect the safety or usefulness of your treatment.
- Tell other doctors or dentists that you are being treated with BRAJACT before you receive any treatment from them.
- If you are still having menstrual periods, BRAJACT may cause your ovaries to stop working, resulting in menopausal symptoms (such as hot flashes) and infertility. Your periods may stop. **This may be permanent**, especially if you are 40 years of age or older. Even if you have stopped having periods after treatment, if you were fertile prior to chemotherapy, you may be able to conceive a pregnancy. Use birth control (but **not** birth control pills) if you could become pregnant, even if you have stopped menstruating because of chemotherapy. Do not breast feed during treatment. Talk to your doctor if you have questions about fertility and birth control after treatment.

Serious Risks of Treatment:

Unexpected and unlikely side effects can occur with any drug treatment. The ones listed below are particularly relevant to your treatment plan:

During treatment:

- **Infection:** The number of white blood cells that help fight infections will be lowered by the chemotherapy drugs, usually starting after about day 7 of each of the 8 cycles. Your blood count is expected to return to normal by day 1 of the next cycle, and will be normal after the 6 months of chemotherapy drugs. If your white blood cell count becomes very low you could get a serious infection. **If you have a fever over 38°C or 100°F, call your cancer doctor immediately (24 hours a day) or go immediately to your nearest Hospital Emergency and tell the doctor you are on chemotherapy.**
- **Increased risk of bleeding:** The number of platelets (special blood cells that help your blood to clot normally after injury) may be lowered by the treatment. They are expected to return to normal by day 1 of next cycle. When the platelet count is low you may be more likely to bruise or bleed. Notify your cancer doctor promptly if you develop large or numerous bruises, or unusual bleeding (eg. nosebleed that won't stop, blood in stool, urine, or sputum). Try to avoid using ASA or ibuprofen, if other pain medications could be used. Talk to your doctor if you feel you need to use one of these medications while on chemotherapy. For patients receiving Warfarin, a modification of the dose may be required based on blood test results.

- **Tissue or vein injury:** Doxorubicin and Paclitaxel can cause tissue injury if they leak out of the vein while being given. Report any sensation of burning or pain to your nurse immediately. Chemotherapy may cause some inflammation and/or scarring in the veins, which may make it difficult to start an IV. Your nurse will help your doctor assess whether a special intravenous device (PICC line or portacath) needs to be considered for your therapy. Pain or tenderness may occur where the needle was placed in your vein. If so, apply cool compresses or soak in cool water for 15-20 minutes several times a day

During or after treatment:

- **Heart Failure:** Rarely, Doxorubicin can have a serious effect on the heart, causing failure of the heart’s pumping action, which results in shortness of breath, fatigue and leg swelling. This can sometimes develop years after treatment. Your treatment may be stopped, and a heart function test done, if there are concerns about your heart function during therapy.
- **Leukemia:** After chemotherapy, there is an increase in the risk of leukemia, a cancer of the white blood cells. The risk is about 1% or less after this type of chemotherapy.
- **Neuropathy:** Paclitaxel can cause you to develop damage to the peripheral nerve endings (the nerves to the hands and feet, and rarely, the face). This can result in feelings of numbness and tingling, or sometimes painful burning sensations. You will need to be careful when handling things that are sharp, hot, or very cold. The majority of the time, these feelings develop after a number of treatments, are not severe, and will resolve fully over a period of months once treatment stops. Infrequently (<5%), these feelings might occur early, might be severe, or might not entirely resolve.

Common chemotherapy side effects and management:

SIDE EFFECT	MANAGEMENT
<i>Nausea and vomiting</i> can occur with AC and you will need to take anti-nausea drugs at the time of the injections and on days 2 and 3 while at home. You will not usually need antinausea drugs for paclitaxel.	<p>You will be given a prescription for anti-nausea drugs to take before your IV treatment and afterwards at home. It is easier to prevent nausea than treat it once it has occurred, so follow directions closely.</p> <ul style="list-style-type: none"> • Drink plenty of liquids. • Eat and drink in small amounts. Try the ideas in <i>Food Choices to Control Nausea</i>*.
Allergic reactions often occur after the administration of Paclitaxel. Signs of an allergic reaction are flushing, rash, itching, dizziness, swelling or breathing problems, or sudden chest or back pain.	<ul style="list-style-type: none"> • Dexamethasone is used to help prevent allergic reactions. You will be given dexamethasone either as a single injection at the time of your treatment. • Your nurse will check your heart rate (pulse) and blood pressure if needed. • Tell your nurse or doctor <i>immediately</i> if you have any sign of an allergic reaction

SIDE EFFECT	MANAGEMENT
<p>Hair loss. Your hair will fall out 2-4 weeks after AC treatment begins. Your scalp may feel tender. You may lose hair on your face and body. Your hair will grow back once your chemotherapy treatments are over and sometimes between treatments. The colour and texture of the new hair growth may be different.</p>	<ul style="list-style-type: none"> • Avoid hair spray, bleaches, dyes and perms. • Apply mineral oil to your scalp to reduce itching. • If you lose your eyelashes and eyebrows, protect your eyes from dust and grit with a broad-rimmed hat and glasses. • Ask your nurse about information on wigs and head coverings.
<p>Fatigue is common especially in the first week after your treatment. As the number of chemotherapy cycles increases, fatigue may get worse or last longer.</p>	<ul style="list-style-type: none"> • Your energy level will improve with time after treatment is completed. • You may obtain a suggestion pamphlet for handling fatigue from nursing staff in your facility
<p>Mouth sores may occur a few days after chemotherapy treatment and may last days or weeks. Mouth sores can occur on the tongue, gums, and the sides of the mouth or in the throat.</p>	<ul style="list-style-type: none"> • Brush your teeth gently after eating and at bedtime with a very soft toothbrush. If your gums bleed, use gauze instead of a brush. Use baking soda instead of toothpaste. • Try baking soda mouth rinses (using 1/4 tsp baking soda in 1 cup warm water) and rinse several times a day. Try ideas in <i>Easy to chew, easy to swallow food ideas*</i>. • Tell your doctor about a sore mouth, as your chemotherapy doses may need to be decreased if mouth sores are severe. Call your doctor if you are having difficulty eating or drinking due to pain.
<p>Diarrhea may occur between treatments.</p>	<p>To help diarrhea:</p> <ul style="list-style-type: none"> • Drink plenty of liquids. • Eat and drink often in small amounts. • You can purchase IMODIUM® over the counter if diarrhea is persistent. Don't take this if you have abdominal pain, blood in your stool, or fever. In that case, consult your doctor promptly for tests. <p>Avoid high fiber foods as outlined in <i>Food ideas to help with diarrhea during chemotherapy</i>.</p>
<p>Burning on urination. AC can cause an irritation of the bladder that result in burning on urination and/or frequency of urination. Urine color may be red after AC for a short while.</p>	<ul style="list-style-type: none"> • Drink more fluids to keep the urine dilute and empty your bladder often.

SIDE EFFECT	MANAGEMENT
<p>Pain affecting joints or muscles may occur for a few days after paclitaxel. After you stop chemotherapy altogether, you may also feel increased joint aching or stiffness for a few months.</p>	<ul style="list-style-type: none"> • Take ibuprofen (e.g., ADVIL®) or acetaminophen (e.g. TYLENOL®) for mild to moderate pain. Contact your cancer doctor if your pain is severe. • Your family doctor can help you to manage symptoms of joint pain after chemotherapy.

*Please ask your chemotherapy nurse, pharmacist or dietician for a copy.

If you experience symptoms or changes in your body that have not been described above but worry you, or if any symptoms are severe, contact

_____ at telephone number _____