

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: BRAJPNCT

(Page 1 of 2)

KEMINDEK: F	lease ensure drug allergies and previous bleomycin are documented on the A	llergy & Alert Form	
DATE:	To be given: Cycle #:		
Date of Previou	s Cycle:		
Number of DC	CEtaxel doses completed to date:		
Number of tra	stuzumab doses completed to date:		
	ment week(s)		
	Platelets day of treatment		
	ith doses as written if within 96 hours ANC <u>greater than or equal to</u> 1.5 x 10 ⁹ /L, PI I x 10 ⁹ /L, Bilirubin <u>less than or equal to</u> 1.5 x ULN, AST or ALT <u>less than or eq</u>		
	on for: Hematology Other Toxicity		
Proceed with	reatment based on blood work from		
PREMEDICA	FIONS: Patient to take own supply. RN/Pharmacist to confirm		
dexamethaso	e 🗌 8 mg or 🗌 12 mg (select one) PO 30 to 60 minutes prior to CARBOplatin		
AND select	ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin		
ONE of the following:	aprepitant 125 mg PO 30 to 60 minutes prior to CARBOplatin, and		
ionowing.	ondansetron 8 mg PO 30 to 60 minutes prior to CARBOplatin		
	netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to CARBOplatin		
If additional an	iemetic required:		
☐ OLANZapiı	e 🗌 2.5 mg or 🔲 5 mg or 🔲 10 mg (select one) PO 30 to 60 minutes prior to CAF	RBOplatin	
Other:			
	Have Hypersensitivity Reaction Tray and Protocol Available		
CHEMOTHE	RAPY: (Note – continued over 2 pages)		
☐ Patients wh	have received only ONE cycle of trastuzumab previously		
trastuzumab 6	mg/kg x kg = mg IV in NS 250 mL over 1 hour. Observe for 30 r	ninutes post-infusion.	
	o select trastuzumab brand as per Provincial Systemic Therapy Policy III-190		
Drug	Brand (Pharmacist to complete. Please print.) Pharmacist Initial a	nd Date	
	imab		
trastuz	<u> </u>		
trastuz	AB (ABRAXANE) 260 mg/m² x BSA = mg		
trastuz	,		
trastuz	B (ABRAXANE) 260 mg/m² x BSA =mg odification:mg/m² x BSA =mg ninutes (in empty sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing wi	ith 15 micron filter)	
PACLitaxel NA Dose M IV over 30	odification:mg/m² x BSA =mg ninutes (in empty sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing wi	ith 15 micron filter)	
PACLitaxel NA Dose M IV over 30 CARBOplatin	odification:mg/m² x BSA =mg ninutes (in empty sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the control of the	ith 15 micron filter)	
trastuz PACLitaxel NA Dose M IV over 30 CARBOplatin Dose M	odification:mg/m² x BSA =mg ninutes (in empty sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing wi	ith 15 micron filter)	
trastuz PACLitaxel NA Dose M IV over 30 CARBOplatin Dose M	odification:mg/m² x BSA =mg ninutes (in empty sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the standard sta	ith 15 micron filter)	
trastuz PACLitaxel NA Dose M IV over 30 CARBOplatin Dose M IV in 100 to	odification:mg/m² x BSA =mg ninutes (in empty sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or n		
trastuz PACLitaxel NA Dose M IV over 30 CARBOplatin Dose M	odification:mg/m² x BSA =mg ninutes (in empty sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with the sterile PVC, non-PVC or n	ith 15 micron filter)	



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: BRAJPNCT

(Page 2 of 2)

DATE:	To be given:	Cycle #:		
CHEMOTHERAPY: (Continued)				
	*** SEE PAGE 1 FOR CHEMOTHERAPY (CYCLE 1 ***		
<u>OR</u> ☐ Patients who have	received TWO cycles or more of trastuzumab previous	sly		
trastuzumab 6 mg/kg infusion (not required a	y x kg = mg IV in NS 250 mL over 30 after 3 treatments with no reaction).	0 minutes. Observe fo	or 30 minutes post-	
Pharmacy to select trastuzumab brand as per Provincial Systemic Therapy Policy III-190				
Drug	Brand (Pharmacist to complete. Please print.)	Pharmacist Initial	and Date	
trastuzumab				
PACLitaxel NAB (ABRAXANE) 260 mg/m² x BSA =mg Dose Modification:mg/m² x BSA =mg IV over 30 minutes (in empty sterile PVC, non-PVC or non-DEHP bag and tubing; use tubing with 15 micron filter) CARBOplatin AUC 6 Dose = AUC x (GFR +25) =mg Dose Modification:% =mg IV in 100 to 250 mL NS over 30 minutes. acetaminophen 325 to 650 mg PO PRN for headache and rigors				
RETURN APPOINTMENT ORDERS				
Return in three we	eeks for Doctor and Cycle			
☐ Last Cycle. Retur				
CBC and Diff, Platele MUGA Scan or Echo trastuzumab and upon If clinically indicated: Other tests: Consults: See general orde				
DOCTOR'S SIGNAT	SIGNATURE:			
			UC:	