

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca/terms-of-use and according to acceptable standards of care.

PROTOCOL CODE: LUMMPP

Page 1 of 1

DOCTOR'S ORDERS	Ht	cm	Wt	kg	BSA	m²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form						
	e given:	•		Cycle #:		
Date of Previous Cycle:				•		
Delay treatment week(s) CBC & Diff, Platelets day of treatment May proceed with doses as written if within 96 hours ANC greater than or equal to 1.5 x 10°/L, Platelets greater than or equal to 100 x 10°/L, Creatinine Clearance: greater than or equal to 45 mL/minute (for pemetrexed and CARBOplatin) or greater than or equal to 60 mL/minute (for CISplatin) Dose modification for: Hematology Other Toxicity: Proceed with treatment based on blood work from						
PREMEDICATIONS: Patient to take own su	upply, RN/Phar	macist to co	nfirm			
dexamethasone 8 mg or 12 mg (select one) PO 30 to 60 minutes prior to treatment						
AND select aprepitant 125 mg PO 30 to 60 minutes prior to treatment, and						
ONE of the ondansetron 8 mg PO 30 to 60 minutes prior to treatment						
following: netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to treatment						
ondansetron 8 mg PO 30 to 60 minutes prior to treatment						
If additional antiemetic required: OLANZapine 2.5 mg or 5 mg or 10 mg (select one) PO 30 to 60 minutes prior to treatment Ensure patient is taking folic acid and has had vitamin B12 injection starting at least 7 days prior to first cycle, and to continue while on treatment, until 21 days after last pemetrexed dose. **Have Hypersensitivity Reaction Tray & Protocol Available**						
HYDRATION:						
1000 mL NS over 1 hour prior to CISplatin						
CHEMOTHERAPY: pemetrexed 500 mg/m² x BSA = mg Dose Modification: % = mg/m² x BSA = mg IV in 100 mL NS over 10 minutes (may be given during prehydration) CISplatin 75 mg/m² x BSA = mg Dose Modification: % = mg/m² x BSA = mg						
IV in 500 mL NS, with potassium chloride 20 mEq, magnesium sulfate 1 g and mannitol 30 g over 1 hour						
OR						
RETURN APPOINTMENT ORDERS						
Return in three weeks for Doctor and Cycl						
Last Cycle. Return in week(s).						
CBC & Diff, Platelets, Creatinine, Bili, ALT, CBC & Diff, Platelets weekly during Cycles 1 Vitamin B12 injection required every 9 weeks This patient to receive injection in clinic. Ne Other tests: Consults: See general orders sheet for additional	and 2 s. Patient to obta xt injection due	ain supply.	·			
DOCTOR'S SIGNATURE:				SIG UC:	NATURE	: