



Provincial Health Services Authority

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca/terms-of-use and according to acceptable standards of care.

PROTOCOL CODE: LUPUPE

DOCTOR'S ORDERS			Ht _____ cm	Wt _____ kg	BSA _____ m ²
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form					
DATE: _____		To be given: _____		Cycle #: _____	
Date of Previous Cycle: _____					
<input type="checkbox"/> Delay treatment _____ week(s) <input type="checkbox"/> CBC & Diff, Platelets day of treatment May proceed with doses as written if within 96 hours ANC greater than or equal to 1.5 x 10⁹/L, Platelets greater than or equal to 100 x 10⁹/L, Creatinine Clearance greater than or equal to 60 mL/minute (if using CISplatin) Dose modification for: <input type="checkbox"/> Hematology <input type="checkbox"/> Other Toxicity _____ Proceed with treatment based on blood work from _____					
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm _____. ondansetron 8 mg PO 30 to 60 minutes prior to treatment on Days 1 to 3 dexamethasone <input type="checkbox"/> 8 mg or <input type="checkbox"/> 12 mg (select one) PO 30 to 60 minutes prior to treatment on Days 1 to 3 <input type="checkbox"/> aprepitant 125 mg PO 30 to 60 minutes prior to treatment on Day 1; then 80 mg PO daily on Day 2 and 3 If additional antiemetic required: <input type="checkbox"/> OLANzapine <input type="checkbox"/> 2.5 mg or <input type="checkbox"/> 5 mg or <input type="checkbox"/> 10 mg (select one) PO 30 to 60 minutes prior to treatment <input type="checkbox"/> hydrocortisone 100 mg IV prior to etoposide <input type="checkbox"/> diphenhydrAMINE 50 mg IV prior to etoposide <input type="checkbox"/> Other: _____					
Have Hypersensitivity Reaction Tray and Protocol Available					
CHEMOTHERAPY:					
CISplatin 25 mg/m²/day x BSA = _____ mg <input type="checkbox"/> Dose Modification: _____ % = _____ mg/m ² x BSA = _____ mg IV in 100 to 250 mL NS over 30 minutes x 3 days OR CARBOplatin AUC 5 x (GFR + 25) = _____ mg IV in 100 to 250 mL NS over 30 minutes Day 1 only etoposide 100 mg/m²/day x BSA = _____ mg <input type="checkbox"/> Dose Modification: _____ % = _____ mg/m ² x BSA = _____ mg IV in 250 to 1000 mL (non-DEHP bag) NS over 45 minutes to 1 hour 30 minutes x 3 days (use non-DEHP tubing with 0.2 micron in-line filter)					
STANDING ORDER FOR ETOPOSIDE TOXICITY:					
hydrocortisone 100 mg IV prn / diphenhydrAMINE 50 mg IV prn					
RETURN APPOINTMENT ORDERS					
<input type="checkbox"/> Return in <input type="checkbox"/> three weeks or <input type="checkbox"/> four weeks (select one) for Doctor and Cycle ____ Book chemo x 3 days. <input type="checkbox"/> Last Cycle. Return in _____ week(s).					
CBC & Diff, Platelets, Creatinine prior to each cycle If clinically indicated: <input type="checkbox"/> Bilirubin <input type="checkbox"/> Other tests: <input type="checkbox"/> Consults: <input type="checkbox"/> See general orders sheet for additional requests.					
DOCTOR'S SIGNATURE:				SIGNATURE:	
				UC:	